

HETEROCYCLES, Vol. 61, 2003, pp. 73 - 78

Received, 30th June, 2003, Accepted, 25th August, 2003, Published online, 1st September, 2003

ELECTROCHEMICAL SYNTHESIS OF SPIROISOXAZOLE DERIVATIVES AND ITS APPLICATION TO NATURAL PRODUCTS


Takahisa Ogamino, Yuichi Ishikawa, and Shigeru Nishiyama*

Department of Chemistry, Faculty of Science and Technology, Keio University,
Hiyoshi 3-14-1, Kohoku-ku, Yokohama 223-8522, Japan

Tel / Fax +81-45-566-1717; E-mail: nishiyama@chem.keio.ac.jp


Abstract – Efficient synthesis of the *trans*-spiroisoxazole (**1**) and its unnatural *cis*-isomer (**2**) was accomplished by employing anodic oxidation of **4**, followed by $\text{Zn}(\text{BH}_4)_2$ reduction. This method was applied to an assembly of the carbon framework of zamamistatin (**12**).

A family of marine natural products, carrying a dibromophenylpyruvic acid oxime moiety,¹ is of interest, because of their diverse biological activities, as well as a typical spiroisoxazole framework (**1**), biogenesis of which might involve epoxidation of an aromatic ring, followed by intramolecular nucleophilic attack of an oxime.² In this context, we accomplished the first total synthesis of aerothionins and aerophobin-1³ by the TFA (thallium (III) trifluoroacetate) oxidation of phenol (**4**),⁴ followed by $\text{Zn}(\text{BH}_4)_2$ reduction of spirodienone (**3**). In particular, reduction of the ketone usually provided unnatural *cis*-form (**2**).⁵ Although the thallium oxidation was a reliable method to give the expected spirodienones, the reaction produced a considerable amount of by-products such as dimers or benzofurans; furthermore, excess amounts of the oxidant were required to complete the reaction. This reaction feature should be reconsidered from the viewpoint of recent environmental concerns. Against such background, an improved synthetic methodology of the spiroisoxazoles (type-**3**) was elaborated by employing anodic oxidation of the corresponding phenol derivatives. We disclose herein our research progress involving synthetic attempts toward zamamistatin (**12**).⁶



The phenol (**4**) for the oxidation was synthesized through the Horner-Wadsworth-Emmons reaction (Scheme 1), while our previous investigation included hydrolysis of the azulactone.^{3a,b}

Thus, bromination of **5**, followed by benzylation afforded **6**, which on coupling with an anion of **7** gave enol ether (**8**) in a good overall yield. On exposure to hydroxylamine hydrochloride, **8** underwent simultaneous deprotection and oximation. Finally, selective hydrogenolysis produced the expected substrate (**4**) for electrolysis.



Scheme 1. Reagents: a. i) pyridinium perbromide / Py, 88%; ii) BnBr, K₂CO₃ / DMF, 90%. b. LHMDS, **7**, 96%. c. i) NH₂OH•HCl / MeOH, 91%; ii) H₂, 10% Pd-C / AcOH – dioxane, 85%.


Anodic oxidation of **4** was attempted at the following procedure. *Procedure:* the oxidation was performed under CCE (constant current electrolysis) or CPE (constant potential electrolysis) conditions employing a glassy carbon beaker (200 or 50 mL) as an anode, a platinum wire as a cathode, and

appropriate supporting salts (*ca.* 0.2 mol / L). As can be seen in Table 1, the MeOH - LiClO₄ combinations (entries 1 - 3) provided by-products (**9**, **10**). Formation of dimer (**10**) implied that insufficient potential caused one-electron oxidation to give diaryl ether through a radical intermediate (**11**). Upon using a constant potential at 1.3 V *vs.* SCE, current was used for other oxidation, resulting in recovery of unreacted **4**. The acidic dioxane and acetone were obviously unfit solvents (entries 4, 5), although they contributed to production of oxa-spiro derivatives.⁷ Eventually, the best condition was obtained when **4** (1 mmol / L) was oxidized under CPE conditions in MeCN - *n*Bu₄NClO₄ (entry 7). Successive reduction of **4** with Zn(BH₄)₂ involving a work-up procedure with MgSO₄, afforded corresponding *trans* and *cis* alcohols (**1**, **2**) in 41 and 39% yields, respectively, which were twice higher than those reported previously.³ In addition, exposure of unnecessary **2** to Et₂NH effected β -elimination to give **4**, which could be repeatedly submitted to the oxidation.


Table 1. Anodic Oxidation of Dibromophenylpyruvic Acid Oxime (**4**)

entry	solvent	supporting salt	potential (V <i>vs.</i> SCE)	current (mA) ^b	conc. of 4 (mmol/L)	yields (%)			
						3	9	10	4
1	MeOH	LiClO ₄	0.9-1.3	10	3	11	8	41	-
2	MeOH	LiClO ₄	0.9-1.3	10	1	18	12	14	-
3	MeOH	LiClO ₄	1.3	80-5	3	19	17	14	20
4	a	LiClO ₄	1.5-1.7	50	3	no reaction			
5	acetone	<i>n</i> Bu ₄ NClO ₄	1.6	50-9	3	multi spots			
6	MeCN	<i>n</i> Bu ₄ NClO ₄	1.6	50-5	3	49	-	-	7
7	MeCN	<i>n</i> Bu ₄ NClO ₄	1.6	20-5	1	68	-	-	-

a. 10% HClO₄ / dioxane. b. Equivalent of current: 2 F / mol.



In the next stage, our attention turned to application of this electrochemical methodology to synthesis of natural products. A literature survey of dibromophenylpyruvic acid derivatives of marine origin indicated zamamistatin (**12**), isolated from the Okinawan sponge *Pseudoceratina purpurea*.⁶ The unprecedented bisspirosisoxazole-structure possessed antibacterial activity against *Rhodospirillum salexigens* SCRC 113, which implied **12** would be a candidate of an inhibitor of biofouling.



Scheme 2. Reagents: a. TBSOTf, 2,6-lutidine. b. LiBH₄, 78% in two steps. c. DMSO, SO₃•Py, Et₃N, 90%. d. i) dimethyl phosphite, Et₃N; ii) TBSCl, Imd., 89%. e. LHMDs / THF, then **6**, 47%. f. i) TBAF; ii) NH₂OH•HCl, 98% in two steps. g. H₂, Pd black / AcOH - dioxane, 54%. h. CPE, 1.6 V vs SCE, 2 F / mol, *n*Bu₄NClO₄ / MeCN, 35%. i. Zn(BH₄)₂ / CH₂Cl₂, **21a**: 26%, **21b**: 23%. j. (S)-MTPACl, DMAP, Py, **22**: 45%, **23**: 43%, **24**: 78%.

After protection of an allyl alcohol of **1** as a siloxy ether (**13**), LiBH₄ reduction provided alcohol (**14**) in 78% yield from **1** (Scheme 2). Compound (**14**) was oxidized to give aldehyde (**15**), which was coupled with dimethyl phosphite under basic conditions, followed by protection of an alcohol generated, leading to the Horner-Wadsworth-Emmons reagent (**16**). Coupling of an anion of **16** with **6** provided **17** in 47% yield. Successively, **17** was converted through **18**⁸ into oxime-phenol (**19**), which on anodic oxidation provided the bispiroisoxazoline derivative (**20**), as a diastereomeric mixture. Zn(BH₄)₂ reduction of **20** gave **21a**⁹ and **21b**⁹ in 26 and 23% yields, respectively. The stereochemistry of **21a** and **21b** could be distinguished by conversion into their (*S*)-MTPA esters: **21a** gave a separable mixture of **22** and **23**, whereas **21b** provided **24** as the sole product. Since zamamistatin (**12**) has a C₂ symmetry, introduction of an optically active MTPA residue to racemic **21a**, would produce a mixture of diastereomers, while *meso*-compound (**21b**) gave *meso*-product (**24**). Accordingly, **21a** should possess the same stereochemistry of the two spiro-centers as zamamistatin (**12**).¹⁰

In conclusion, we have accomplished an improved synthetic method of the spiroisoxazole derivatives (**1**, **2**) by using anodic oxidation. This method provided access to the assembly of the zamamistatin-framework. Further investigation towards the natural zamamistatin molecule and related natural products carrying dibromophenylpyruvic acid oxime is in progress.

ACKNOWLEDGMENTS

This work was supported by Grant-in-Aid for the 21st Century COE program "KEIO Life Conjugate Chemistry", as well as Scientific Research C (15510181), from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

REFERENCES AND NOTES

1. J. D. Faulkner, *Nat. Prod. Rep.*, 2002, **19**, 1. Many references are cited therein.
2. K. T. Okamoto and J. Clardy, *Tetrahedron Lett.*, 1987, **28**, 4969.
3. (a) S. Nishiyama and S. Yamamura, *Tetrahedron Lett.*, 1983, **24**, 3351. (b) S. Nishiyama and S. Yamamura, *Bull. Chem. Soc. Jpn.*, 1986, **58**, 3453. (c) Other arothionin synthesis: H. H.

- Wasserman and J. Wang, *J. Org. Chem.*, 1998, **63**, 5581.
4. Related oxidation: (a) M. Kaçan, D. Koyuncu, and A. Mckillop, *J. Chem. Soc., Perkin Trans. I*, **1993**, 1771. (b) H. Hara, T. Inoue, H. Nakamura, M. Endoh, and O. Hoshino, *Tetrahedron Lett.*, 1992, **33**, 6491. (c) M. Murakata, k. Yamada, and O. Hoshino, *J. Chem. Soc., Chem. Commun.*, **1994**, 443. (d) M. Murakata, K. Yamada, and O. Hoshino, *Tetrahedron*, 1996, **52**, 14713. (e) M. Murakata, M. Tamura, and O. Hoshino, *J. Org. Chem.*, 1997, **62**, 4428.
 5. A. R. Forrester, R. H. Thomson, and S.-O. Woo, *Liebigs Ann. Chem.*, **1978**, 66. Wasserman *et al.* reported that the *trans*-product was obtained in 25% yield under NaBH₃CN / TFA conditions.^{3c}
 6. N. Takada, R. Watanabe, K. Suenaga, K. Yamada, K. Ueda, M. Kita, and D. Uemura, *Tetrahedron Lett.*, 2001 **42**, 5265.
 7. F. Doi, T. Ogamino, T. Sugai, and S. Nishiyama, *Synlett*, **2003**, 411.
 8. The TBAF treatment effected a high-yield conversion of the TBS-enol ether into the oxime. Upon a direct use of NH₂OH•HCl as in the case of **8**, **18** was obtained in 39% yield.
 9. **21a**: IR (film) 3434 cm⁻¹; NMR δ_{H} (CDCl₃) 2.29 (2H, d, *J*= 8.3 Hz), 3.03 (2H, d, *J*= 17.6 Hz), 3.76 (6H, s), 4.00 (2H, d, *J*= 17.6 Hz), 4.44 (2H, d, *J*= 8.3 Hz), 6.33 (2H, s); δ_{C} (CDCl₃) 38.6, 60.2, 73.9, 91.0, 112.3, 121.1, 131.0, 148.0, 150.9. HRMS *m/z* (M⁺-OH) Calcd for C₁₈H₁₅N₂O₅⁷⁹Br₄, 654.7715. Found: 654.7755. **21b**: IR (film) 3436 cm⁻¹; NMR δ_{H} (acetone-d₆) 3.23 (2H, d, *J*= 17.6 Hz), 3.72 (6H, s), 3.87 (2H, d, *J*= 17.6 Hz), 4.15 (2H, d, *J*= 8.3 Hz), 5.44 (2H, d, *J*= 8.3Hz), 6.51 (2H, s); δ_{C} (acetone-d₆) 39.7, 60.2, 75.1, 91.3, 113.7, 122.1, 132.0, 148.6, 151.9. HRMS *m/z* (M⁺) Calcd for C₁₈H₁₆N₂O₆⁷⁹Br₄, 671.7682. Found: 671.7712.
 10. Unfortunately, efforts to convert into natural product (**12**) have been unsuccessful up to now, and continuous attempts are still in progress.