

HETEROCYCLES, Vol. 84, No. 2, 2012, pp. 1045 - 1056. © 2012 The Japan Institute of Heterocyclic Chemistry  
 Received, 14th July, 2011, Accepted, 15th August, 2011, Published online, 16th August, 2011  
 DOI: 10.3987/COM-11-S(P)82

## CHIBA-G-CATALYZED INTRAMOLECULAR OXO-MICHAEL ADDITION: SYNTHETIC APPROACHES TO VITAMINE E SKELETON

Sayaka Tokunou, Waka Nakanishi, Natsuko Kagawa, Takuya Kumamoto,  
 and Tsutomu Ishikawa\*

Graduate School of Pharmaceutical Sciences, Chiba University, 1-33 Yayoi,  
 Inage, Chiba 263-8522, Japan

E-mail: benti@faculy.chiba-u.jp

**Abstract** – A chroman skeleton with quaternary carbon chiral center, leadable to vitamin E after manipulation, was constructed through 6-*exo-trig* type intramolecular oxo-Michael addition in up to 44% yield with 81% *ee* when a phenol with (*Z*)- $\alpha,\beta$ -unsaturated ester at *ortho* position was reacted in the presence of a guanidine-type organocatalyst, Chiba-G [(-)-(4*R*,5*R*)-2-[(*S*)-1-hydroxymethyl-2-phenylethyl]imino-1,3-dimethyl-4,5-diphenylimidazolidine (or the enantiomer)].

### INTRODUCTION

Development of enantioselective construction of 2,2-disubstituted chroman skeletons is an important issue in the synthesis of biologically active natural products, such as vitamin E (**1**)<sup>1</sup> with lipophilic antioxidant activity and *trans*- $\delta$ -tocotrienoloic acid (**2**)<sup>2</sup> with antibacterial activity, for the purpose of further therapeutic modification (Figure 1).





Figure 1. Structures of vitamin E (**1**), *trans*- $\delta$ -tocotrienoloic acid (**2**) and (-)-Chiba-G

Hayashi,<sup>3</sup> Achiwa,<sup>4</sup> Trost,<sup>5</sup> and Tietze<sup>6</sup> and their co-workers have reported the palladium-catalyzed asymmetric synthesis of 2,2-disubstituted chiral chromans. However, although much attention has been paid for the use of organocatalyst instead of metal catalyst in asymmetric synthesis,<sup>7</sup> there had been no reports on the construction of 2,2-disubstituted chiral chroman skeletons by using organocatalyst. We have uncovered potential functionality of guanidine compounds as chiral auxiliaries<sup>8</sup> and recently reported the model asymmetric construction of 2,2-disubstituted chroman skeleton from phenols with tri-substituted  $\alpha,\beta$ -unsaturated esters by 6-*exo-trig* type intramolecular oxo-Michael addition (IOMA) catalyzed by chiral guanidines.<sup>9</sup> In this paper we present the synthetic application of this methodology using a backbone guanidine catalyst in the model reaction, Chiba-G [(-)-(4*R*,5*R*)-2-[(*S*)-1-hydroxymethyl-2-phenylethyl]imino-1,3-dimethyl-4,5-diphenylimidazolidine (or its enantiomer)] (Figure 1), to the construction of vitamin E core.


## RESULTS AND DISCUSSION

**Trials for IOMA of 2-hydroxy-5-methoxy-3,4,6-trimethylphenyl vinyl ketone 9: 6-*endo-trig* type reaction.** We have developed 6-*endo-trig* type IOMA catalyzed by quinine for the preparation of 2,3-disubstituted chroman-4-one system and applied the method to the enantioselective synthesis of anti-HIV-1 active coumarins such as (+)-calanolide A.<sup>10</sup> Thus, at first, we examined Chiba-G-catalyzed 6-*endo-trig* type IOMA of 2-hydroxy-5-methoxy-3,4,6-trimethylphenyl vinyl ketone **9**, which was prepared as shown in Scheme 1. Reaction of a MOM-protected benzaldehyde **5**, which was derived from a commercially available 2,3,6-trimethylhydroquinone (**3**) by conventional methods in 4 steps, with 6-benzyloxyhex-1-yne afforded an alkynyl alcohol **6** in 67% yield. Oxidation of **6** with MnO<sub>2</sub> followed by treatment with



Scheme 1. Preparation of substrate **9** for 6-*endo-trig* type IOMA

Me<sub>2</sub>CuLi yielded an alkenyl ketone **8**. After the repeated chromatographic separation of the *E/Z*-isomers of **8**, removal of the MOM-deprotection was examined under various conditions. The use of NaHSO<sub>4</sub>-SiO<sub>2</sub> complex<sup>11</sup> successfully provided a targeted vinyl ketone **9** in high yield; however, a partial isomerization was observed during the deprotection [*E* : *Z* = 1 : 0.3 from *E*-**7**; *E* : *Z* = 0.25 : 1 from *Z*-**7**]. The results of separate IOMA on (*E*)- and (*Z*)-enriched 2-hydroxy-5-methoxy-3,4,6-trimethylphenyl vinyl ketone **9** were summarized in Table 1. No reactions occurred in the presence of a tertiary amine such as quinine or quinidine (runs 1 and 2), while Chiba-G worked as a catalyst; but both chemical yield and enantioselectivity of cyclized product **10** were low in the use of CHCl<sub>3</sub> as solvent (runs 3 and 4) or without solvent (run 6). Reaction rate was greatly accelerated when reaction was carried out in MeOH; however, no asymmetric induction was observed (run 5). The recovered starting vinyl ketone **9** was found to be an (*E*)-enriched mixture, even from a (*Z*)-enriched **9**, suggesting a possible retro-IOMA. Thus, the 6-*endo-trig* type IOMA of a vinyl ketone **9** resulted in unfruitful cyclization.


Table 1. 6-*endo-trig* type IOMA of a vinyl ketone **9**

run	<b>9</b> ( <i>E</i> : <i>Z</i> )	base / solvent	time (d)	<b>10</b>		<b>9</b>	
				yield (%)	ee(%)	yield (%)	( <i>E</i> : <i>Z</i> )
1	0.2 : 1	quinine / CHCl <sub>3</sub>	4	0	-	76	1 : 0.8
2	0.2 : 1	quinidine / CHCl <sub>3</sub>	4	0	-	73	1 : 0.7
3	1 : 0.3	(-)-Chiba-G / CHCl <sub>3</sub>	1	29	4	51	1 : 0.5
4	0.25 : 1	(-)-Chiba-G / CHCl <sub>3</sub>	1	37	4	33	1 : 0.5
5	0.4 : 1	(-)-Chiba-G / MeOH	1	95	0	0	-
6	0.4 : 1	(-)-Chiba-G / none	6	20	0	91	1 : 0.4

**Trials for IOMA of methyl 5-(2-hydroxy-5-methoxy-3,4,6-trimethylphenyl)-3-methylpent-2-enoate (**14**): 6-*exo-trig* type reaction.** Next, we turned our attention to 6-*exo-trig* type reaction and methyl 5-(2-hydroxy-5-methoxy-3,4,6-trimethylphenyl)-3-methylpent-2-enoate (**14**) was chosen as a substrate for the IOMA. The ester **14** was prepared from the MOM-protected benzaldehyde **5**, according to the procedure adopted in our model examination<sup>9</sup> (Scheme 2). Four-successive reactions of aldol condensation with acetone, catalytic hydrogenation, Horner-Emmons-Wadsworth reaction with methyl *p,p*-bis(2,2,2-trifluoroethyl)phosphonoacetate, and the MOM-deprotection afforded (*E*)-methyl ester *E*-**14** and (*Z*)-methyl ester *Z*-**14**.


Smooth cyclization of both (*E*)- and (*Z*)-isomers was, as expected, observed in the tetramethylguanidine (TMG)-catalyzed reactions (Table 2, runs 1 and 2). Similarly, Chiba-G worked well as a catalyst in the

reaction of (*E*)-substrate **E-14** to give a chroman product **15** in 81% yield, but selectivity was not satisfactory (30% ee) (run 3). On the other hand, acceptable enantioselectivity (81% ee) was obtained, even moderate chemical yield (44%), when (*Z*)-substrate **Z-14** was subjected to the IOMA (run 4). This



Scheme 2. Preparation of substrate **14** for 6-*exo-trig* type IOMA

Table 2. Guanidine-catalyzed 6-*exo-trig* type IOMA of an unsaturated ester **14**




run	<b>14</b>	guanidine / solvent	temp. (°C)	time	<b>15</b>	
					yield (%)	ee (%)
1	<i>E</i>	TMG / CHCl <sub>3</sub>	rt	7 h	77	-
2	<i>Z</i>	TMG / CHCl <sub>3</sub>	rt	3 d	93	-
3	<i>E</i>	(-)-Chiba-G / CHCl <sub>3</sub>	rt	4 d	81	30 ( <i>S</i> )
4	<i>Z</i>	(-)-Chiba-G / CHCl <sub>3</sub>	rt	4 d	44 <sup>a</sup>	81 ( <i>R</i> )
5	<i>Z</i>	(-)-Chiba-G / CHCl <sub>3</sub>	50	2 d	62 <sup>a</sup>	74 ( <i>R</i> )
6	<i>Z</i>	(-)-Chiba-G / PhMe	rt	4 d	23 <sup>a</sup>	58 ( <i>R</i> )
7	<i>Z</i>	(-)-Chiba-G / THF	rt	4 d	27 <sup>a</sup>	63 ( <i>R</i> )
8	<i>Z</i>	(-)-Chiba-G / acetone	50	2 d	14 <sup>a</sup>	62 ( <i>R</i> )
9	<i>Z</i>	(-)-Chiba-G / DMF	rt	4 d	38 <sup>a</sup>	3 ( <i>R</i> )

<sup>a</sup> Estimated by <sup>1</sup>H NMR

reaction was tolerant to temperature and, interestingly, the increment of chemical yield (62%) with slight loss of enantioselectivity (81%  $\rightarrow$  74% ee) was observed (run 5). Unfortunately, both chemical yield and enantioselectivity in the IOMA of **Z-14** were not improved in the screening of solvents (runs 6-9); however, no occurrence of a retro-path in these reactions was suggested by the recovery of the starting **Z-14**.

Thus, it was found that chloroform was the best solvent among solvents examined for the 6-*exo-trig* type IOMA of **Z-14** (run 4 in Table 2). We had observed the rate acceleration of 6-*endo-trig* type IOMA in calanolide A synthesis when chlorobenzene was used as a solvent,<sup>10</sup> Next, (-)-Chiba-G-catalyzed IOMA of **Z-14** in a chlorinated solvent was examined (Table 3). In general, high enantioselectivity, especially 87% ee in the use of dichloromethane (run 4), was obtained, but chemical yields were not improved in spite of their permittivity.


Table 3. (-)-Chiba-G-catalyzed IOMA of **Z-14** in a chlorinated solvent



run	solvent	relative permittivity ( $\epsilon$ )	<b>15</b>	
			yield (%)	ee (%)
1	CCl <sub>4</sub>	2.30	7	79
2	PhCl	5.74	11	82
3 <sup>a</sup>	CHCl <sub>3</sub>	4.89	44	81
4	CH <sub>2</sub> Cl <sub>2</sub>	9.02	39	87
5	(CH <sub>2</sub> Cl <sub>2</sub> ) <sub>2</sub>	10.74	25	84

<sup>a</sup> The data of run 4 in Table 2.

Absolute configuration of the cyclization product **15** was determined by comparison with a known (*S*)-(+)-vinylous chroman *S*-(+)-**17**  $\{[\alpha]_D +15.4 (\text{CHCl}_3)\}$ .<sup>6</sup> DIBAH-reduction of (+)-**15**  $\{[\alpha]_D +2.8 (\text{CHCl}_3)\}$  with 77% ee, derived from the (-)-Chiba-G-catalyzed IOMA of **Z-14**, followed by Wittig reaction afforded (-)-vinylous chroman *R*-(-)-**17**  $\{[\alpha]_D -10.8 (\text{CHCl}_3)\}$ , the ee of which was estimated to be 68% (Scheme 3). These facts indicated that asymmetric induction leading *R*-configuration was controlled in the (-)-Chiba-G-catalyzed IOMA of **Z-14**, with accordance to the results of our model IOMA.<sup>9</sup>



Scheme 3. Determination of the absolute configuration of (+)-chroman (+)-**15**

## CONCLUSION

We examined 6-*endo-trig* and 6-*exo-trig* type IOMAs for the asymmetric construction of vitamin E core carrying a 2,2-disubstituted chroman skeleton and found that the latter 6-*exo-trig* type cyclization could be effectively catalyzed by Chiba-G when (*Z*)-unsaturated ester was used as a substrate.

## EXPERIMENTAL

**General.** Melting points were determined on a micro melting point hot-stage instrument Yanagimoto MP-SI and are uncorrected. IR spectra were recorded with ATR on a JASCO FT/IR-300E spectrometer. Specific rotation,  $[\alpha]_D$ , was recorded on a JASCO DIP-140 polarimeter. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded with JEOL JNM ECP 400 spectrometer in CDCl<sub>3</sub>. HREIMS was performed on JASCO MS-GCMATE spectrometer. For column chromatography silica gel 60 or 60N (spherical, 70-230 mesh, Kanto) and for flash chromatography silica gel (230-400 mesh, Merck) were used.

**Compound 6.** To a solution of 6-benzyloxy-1-hexyne (3.76 g, 20.0 mmol) in THF (40 mL) was added a 1.57 M solution of *n*BuLi in hexane (13.5 mL, 21.2 mmol) at 0 °C over 5 min under Ar, and the mixture was stirred at the same temperature for 1 h and then cooled to -42 °C. To the cooled mixture was added a solution of **5** (4.07 g, 17.1 mmol) in THF (30 mL) over 10 min, and the whole was stirred at -42 °C for 1 h, quenched with sat. NaHCO<sub>3</sub> (50 mL), and extracted with AcOEt (50 mL x 3). The combined organic solutions were washed with brine (50 mL), dried (MgSO<sub>4</sub>), and evaporated. Column chromatography of the residue (hexane : AcOEt = 6 : 1) afforded **6** (4.86 g, 67%) as a colorless oil; IR  $\nu_{\max}$ : 3422, 2361 cm<sup>-1</sup>; <sup>1</sup>H NMR  $\delta$ : 1.58-1.64, 1.67-1.72 (each 2H, m, CH<sub>2</sub>), 2.16, 2.19, 2.44 (each 3H, s, Me), 2.25 (2H, dt, *J* = 7.0, 2.1 Hz, CH<sub>2</sub>), 3.47 (2H, t, *J* = 6.3 Hz, CH<sub>2</sub>), 3.57 (1H, d, *J* = 7.4 Hz, OH, exchangeable), 3.635, 3.943 (each 3H, s OMe), 4.48 (2H, s, OCH<sub>2</sub>O), 4.94, 5.01 (each 1H, d, *J* = 5.7 Hz, OCH<sub>2</sub>Ph), 5.85 (1H, dt, *J* =

7.4, 2.1 Hz, CH), 7.28-7.36 (5H, m, ArH);  $^{13}\text{C}$  NMR  $\delta$ : 12.1, 12.6, 13.1, 18.4, 25.0, 28.5, 57.3, 58.5, 59.6, 69.4, 72.4, 80.5, 85.1, 99.8, 127.1, 127.2, 127.6, 128.0, 130.5, 131.2, 138.2, 149.2, 153.5; HREIMS:  $m/z$  426.2409 (calcd for  $\text{C}_{26}\text{H}_{34}\text{O}_5$ : 426.2406).

**Compound 7.** A mixture of **6** (0.392 g, 0.92 mmol) and  $\text{MnO}_2$  (0.804 g, 9.25 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was stirred at rt for 10 h and filtered through celtie pad. Evaporation of the filtrate gave **7** (0.358 g, 92%) as a yellow oil; IR  $\nu_{\text{max}}$ : 2205, 1651  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$ : 1.69-1.72 (4H, m,  $\text{CH}_2 \times 2$ ), 2.18, 2.21, 2.25 (each 3H, s, Me), 2.43 (2H, dif. t,  $J = 6.6$  Hz,  $\text{CH}_2$ ), 3.48 (2H, dif. t,  $J = 5.8$  Hz,  $\text{CH}_2$ ), 3.54, 3.65 (each 3H, s OMe), 4.48 (2H, s,  $\text{OCH}_2\text{O}$ ), 4.89 (2H, s,  $\text{OCH}_2\text{Ph}$ ), 7.28-7.36 (5H, m, ArH);  $^{13}\text{C}$  NMR  $\delta$ : 12.1, 12.6, 18.5, 24.1, 28.4, 57.1, 59.5, 69.0, 72.3, 82.5, 95.7, 100.1, 125.4, 127.0, 127.8, 129.0, 132.8, 133.5, 138.0, 148.2, 152.9, 181.0; HREIMS:  $m/z$  424.2262 (calcd for  $\text{C}_{26}\text{H}_{32}\text{O}_5$ : 424.2250).

**Compound 8.** A 0.83 M solution of MeLi in  $\text{Et}_2\text{O}$  (24 mL, 19.9 mmol) was added to a mixture of CuI (1.87 g, 9.82 mmol) in THF (40 mL) at  $-40$  °C over 10 min, and the whole was stirred at the same temperature for 15 min. After addition of a solution of **7** (1.81 g, 4.26 mmol) in THF (20 mL) over 20 min at  $-40$  °C the mixture was stirred at the same temperature for 20 min, poured into ice-water (150 mL), and extracted with AcOEt (100 mL  $\times$  3). The combined organic solutions were washed with brine (50 mL), dried ( $\text{MgSO}_4$ ), and evaporated. Column chromatography of the residue (hexane : AcOEt = 15 : 1) afforded a 1 : 0.6 mixture of *E*- and *Z*-**8** (1.87 g, 100%) as a pale yellow oil, which could be separated to be each isomer by repeated chromatographies; IR  $\nu_{\text{max}}$ : 1669  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$ : For (*E*)-isomer: 1.60-1.61 (4H, m,  $\text{CH}_2 \times 2$ ), 2.13, 2.15, 2.19, 2.21 (each 3H, s, Me), 2.18-2.19 (2H, m,  $\text{CH}_2$ ), 3.46-3.48 (2H, m,  $\text{CH}_2$ ), 3.46, 3.65 (each 3H, s OMe), 4.49 (2H, s,  $\text{OCH}_2\text{O}$ ), 4.83 (2H, s,  $\text{OCH}_2\text{Ph}$ ), 6.26 (1H, s, CH), 7.31-7.35 (5H, m, ArH); For (*Z*)-isomer: 1.58-1.70 (4H, m,  $\text{CH}_2 \times 2$ ), 1.91, 2.13, 2.18, 2.20 (each 3H, s, Me), 2.64 (2H, dif. t,  $J = 8.1$  Hz,  $\text{CH}_2$ ), 3.46, 3.64 (each 3H, s OMe), 3.51 (2H, dif. t,  $J = 8.1$  Hz,  $\text{CH}_2$ ), 4.52 (2H, s,  $\text{OCH}_2\text{O}$ ), 4.83 (2H, s,  $\text{OCH}_2\text{Ph}$ ), 6.26 (1H, s, CH), 7.28-7.35 (5H, m, ArH);  $^{13}\text{C}$  NMR  $\delta$ : For (*E*)-isomer: 12.44, 12.89, 12.98, 19.3, 24.2, 29.2, 41.0, 57.4, 59.98, 69.8, 72.78, 100.3, 125.14, 126.0, 127.42, 127.496, 128.24, 129.05, 136.10, 138.4, 147.51, 153.323, 159.0, 196.4; For (*Z*)-isomer: 12.43, 12.88, 12.97, 24.8, 25.5, 29.7, 33.5, 57.5, 59.96, 70.0, 72.75, 100.4, 125.12, 126.37, 127.35, 127.504, 128.22, 129.03, 136.13, 138.6, 147.49, 153.316, 159.9, 195.8; HREIMS:  $m/z$  440.2572 (calcd for  $\text{C}_{27}\text{H}_{36}\text{O}_5$ : 440.2563).

**Compound 9 from E-8.** A mixture of *E*-**8** (25.3 mg, 57.4  $\mu\text{mol}$ ) and  $\text{NaHSO}_4 \cdot \text{SiO}_2$  (12.0 mg) in  $\text{CH}_2\text{Cl}_2$  (0.5 mL) was stirred at rt for 1 h and filtered through celite pad. After evaporation of the filtrate column chromatography of the residue (hexane : AcOEt = 15 : 1) gave a 1 : 0.3 mixture of *E*- and *Z*-**9** (21.8 mg, 96%) as a yellow oil; IR  $\nu_{\text{max}}$ : 3734, 1633  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$ : For (*E*)-isomer: 1.61-1.71 (4H, m,

CH<sub>2</sub> x 2), 2.16-2.18 (6H, br, Me x 2), 2.19-2.27 (2H, m, CH<sub>2</sub>), 2.24, 2.39 (each 3H, s, Me), 3.48-3.53 (2H, m, CH<sub>2</sub>), 3.65 (3H, s OMe), 4.51 (2H, s, OCH<sub>2</sub>Ph), 6.34-6.35 (1H, m, CH), 7.25-7.34 (5H, m, ArH), 11.26 (1H, s, OH, exchangeable); For (*Z*)-isomer: 1.61-1.71 (4H, m, CH<sub>2</sub> x 2), 1.96 (3H, d, *J* = 1.3 Hz, Me), 2.16, 2.24, 2.39 (each 3H, s, Me), 2.61 (2H, dif. t, *J* = 7.6 Hz, CH<sub>2</sub>), 3.48-3.53 (2H, m, CH<sub>2</sub>), 3.65 (3H, s OMe), 4.51 (2H, s, OCH<sub>2</sub>Ph), 6.32-6.33 (1H, m, CH), 7.25-7.34 (5H, m, ArH), 11.30 (1H, s, OH, exchangeable); <sup>13</sup>C NMR δ: For (*E*)-isomer: 11.7, 13.3, 15.8, 19.7, 24.1, 29.4, 40.9, 60.1, 69.8, 72.9, 121.5, 123.7, 126.7, 127.49, 127.53, 127.58, 128.30, 136.8, 138.4, 149.6, 154.9, 157.9, 198.0; For (*Z*)-isomer: 11.7, 13.3, 15.7, 24.8, 25.3, 29.7, 33.7, 60.1, 70.0, 72.8, 121.4, 123.7, 126.7, 127.4, 127.55, 127.58, 128.27, 136.8, 138.6, 149.6, 155.0, 158.0, 197.7; HREIMS: *m/z* 396.2297 (calcd for C<sub>25</sub>H<sub>32</sub>O<sub>4</sub>: 396.2300).

**IOMA of 9 (Table 1, run 3): 2-(4-Benzyloxybutyl)-6-methoxy-2,5,7,8-tetramethylbenzo[*b*]pyran-4(3*H*)-one (10).** A solution of an (*E*)-major 9 (22.8 mg, 57.5 μmol) and (-)-Chiba-G (4.6 mg, 11.5 μmol) in CHCl<sub>3</sub> (0.5 mL) was stirred at rt and then under reflux (total 1 day). After evaporation of the solvent, flash chromatography of the residue (hexane : AcOEt = 15 : 1) afforded 10 (6.7 mg, 29%) as a yellow oil together with the recovery of the starting 9 (*E* : *Z* = 1 : 0.5) (11.6 mg, 51%); IR ν<sub>max</sub>: 1678 cm<sup>-1</sup>; <sup>1</sup>H NMR δ: 1.35 (3H, s, Me), 1.46-1.68 (5H, m, CH<sub>2</sub> x 2, CH<sub>2</sub> x 1/2), 1.78 (1H, ddd, *J* = 13.6, 11.6, 5.3 Hz, CH<sub>2</sub> x 1/2), 2.12, 2.24, 2.54 (each 3H, s, Me), 2.58, 2.72 (each 1H, d, *J* = 15.7 Hz, CH<sub>2</sub>), 3.47 (2H, t, *J* = 6.2 Hz, CH<sub>2</sub>), 3.62 (3H, s, OMe), 4.49 (2H, s, OCH<sub>2</sub>Ph), 7.27-7.36 (5H, m, ArH); <sup>13</sup>C NMR δ: 12.0, 13.6, 13.9, 20.3, 23.7, 29.9, 39.2, 49.2, 60.3, 70.0, 72.9, 79.6, 117.1, 124.1, 127.5, 127.6, 128.3, 129.5, 138.2, 138.5, 150.4, 154.8, 194.7; HREIMS *m/z*: 396.2308 (calcd for C<sub>25</sub>H<sub>32</sub>O<sub>4</sub>: 396.2300); HPLC (CHIRALCEL OD-H, λ = 254 nm, eluent: hexane : *i*PrOH = 100 : 1, flow rate: 1.1 mL/min): *t*<sub>R</sub> for a major 10: 24.7 min (52%), *t*<sub>R</sub> for a minor 10: 29.6 min (48%).

**Compound 11.** A mixture of 5 (0.657 g, 2.76 mmol) and acetone (0.67 mL, 9.12 mmol) in 10% NaOH (2.6 mL, 6.5 mmol) and H<sub>2</sub>O (4 mL) was stirred at 45 °C for 1 day under Ar, quenched with H<sub>2</sub>O (10 mL), and extracted with CHCl<sub>3</sub> (20 mL and then 10 mL x 2). The combined organic solutions were washed with brine (10 mL), dried (MgSO<sub>4</sub>), and evaporated. Column chromatography of the residue (hexane : AcOEt = 10 : 1) afforded 11 (0.645 g, 84%) as colorless solid; IR ν<sub>max</sub>: 1682 cm<sup>-1</sup>; <sup>1</sup>H NMR δ: 2.21, 2.23, 2.31, 2.38 (each 3H, s, Me), 3.56, 3.65 (each 3H, s OMe), 4.84 (2H, s, OCH<sub>2</sub>O), 6.67 (1H, d, *J* = 16.7 Hz, CH), 7.75 (1H, d, *J* = 16.7 Hz, C4-H); <sup>13</sup>C NMR δ: 13.25, 13.32, 13.6, 27.6, 57.9, 60.2, 99.9, 126.4, 128.4, 129.1, 132.1, 132.7, 139.5, 150.9, 153.6, 198.8; Anal. Calcd for C<sub>16</sub>H<sub>22</sub>O<sub>4</sub>: C, 69.04; H, 7.97. Found: C, 69.00; H, 8.07.



**Compound 12.** A mixture of **11** (0.642 g, 2.31 mmol) and 5% Pd/C (0.057 g) in AcOEt (20 mL) was stirred at rt for 2 h under H<sub>2</sub> atmosphere and filtered through celite pad. After evaporation of the filtrate column chromatography of the residue (hexane : AcOEt = 7 : 1) afforded **12** (0.567 g, 88%) as a colorless oil; IR  $\nu_{\max}$ : 1711 cm<sup>-1</sup>; <sup>1</sup>H NMR  $\delta$ : 2.16 (6H, s, Me x 2), 2.18, 2.21 (each 3H, s, Me), 2.64, 2.91 (each 2H, dif. t,  $J$  = 8.1 Hz, CH<sub>2</sub>), 3.58, 3.63 (each 3H, s, OMe), 4.89 (2H, s, OCH<sub>2</sub>O); <sup>13</sup>C NMR  $\delta$ : 12.2, 12.8, 13.6, 21.8, 29.8, 43.6, 57.4, 60.1, 99.9, 127.1, 128.2, 128.6, 131.1, 150.7, 153.4, 208.6; HREIMS  $m/z$ : 280.1671 (calcd for C<sub>16</sub>H<sub>24</sub>O<sub>4</sub>: 280.1674).

**Compound 13.** A 60% mineral oil of NaH (0.308 g, 7.71 mmol) was washed with dry hexane (5 mL) and suspended in THF (6 mL). To the suspension was added methyl *p,p*-bis(2,2,2-trifluoroethyl)-phosphonoacetate (1.3 mL, 6.13 mmol) under ice-cooling over 10 min, and the mixture was stirred at rt for 1 h. To the mixture was added a solution of **12** (0.839 g, 2.99 mmol) in THF (7 mL) over 5 min, and the whole was stirred at rt for 4.5 h, quenched with H<sub>2</sub>O (50 mL), and extracted with AcOEt (10 mL x 3). The combined organic solutions were washed with brine (20 mL), dried (MgSO<sub>4</sub>), and evaporated. Column chromatography of the residue (hexane : AcOEt = 12 : 1) afforded a 1 : 0.8 mixture of *E*- and *Z*-**13** (0.890 g, 89%) as colorless solid; IR  $\nu_{\max}$ : 1716 cm<sup>-1</sup>; <sup>1</sup>H NMR  $\delta$ : For (*E*)-isomer: 2.16, 2.18, 2.23 (each 3H, s, Me), 2.26 (3H, d,  $J$  = 1.3 Hz, Me), 2.27-2.32, 2.79-2.83 (each 2H, m, CH<sub>2</sub>), 3.60, 3.64, 3.71 (each 3H, s, OMe), 4.90 (2H, s, OCH<sub>2</sub>O), 5.77 (1H, m, CH); For (*Z*)-isomer: 1.97 (3H, d,  $J$  = 1.3 Hz, Me), 2.18 (6H, s, Me x 2), 2.32 (3H, s, Me), 2.79-2.83 (4H, m, CH<sub>2</sub> x 2), 3.62, 3.65, 3.68 (each 3H, s, OMe), 4.92 (2H, s, OCH<sub>2</sub>O), 5.69 (1H, m, CH); <sup>13</sup>C NMR  $\delta$ : For (*E*)-isomer: 12.1, 12.8, 13.6, 18.9, 25.3, 40.8, 50.8, 57.4, 60.0, 99.8, 114.8, 126.9, 128.1, 128.6, 131.2, 150.6, 153.3, 160.3, 167.2; For (*Z*)-isomer: 12.0, 12.8, 13.6, 26.0, 26.2, 33.4, 50.7, 57.5, 60.0, 99.8, 115.9, 127.6, 128.0, 128.3, 131.7, 150.4, 153.2, 159.9, 166.5; HREIMS:  $m/z$  336.1921 (calcd for C<sub>19</sub>H<sub>28</sub>O<sub>5</sub>: 336.1937).

**Compound 14.** A mixture of **13** (0.825 g, 2.45 mmol) in MeOH (35 mL) containing 10% HCl (7.8 mL, 21.4 mmol) was heated at 70 °C for 1 h, quenched with H<sub>2</sub>O (60 mL), and extracted with AcOEt (30 mL x 3). The combined organic solutions were washed with brine (20 mL), dried (MgSO<sub>4</sub>), and evaporated. Column chromatography of the residue (hexane : AcOEt = 15 : 1 to 7 : 1) afforded *E*-**14** (0.412 g, 58%) and *Z*-**14** (0.291 g, 41%). (i) *E*-**14**: colorless solids, mp 88 °C; IR  $\nu_{\max}$ : 3375, 1712 cm<sup>-1</sup>; <sup>1</sup>H NMR  $\delta$ : 2.14, 2.20, 2.22 (each 3H, s, Me), 2.25 (3H, d,  $J$  = 1.3 Hz, Me), 2.27-2.31, 2.76-2.80 (each 2H, m, CH<sub>2</sub>), 3.62, 3.70 (each 3H, s, OMe), 4.55 (1H, s, OH, exchangeable), 5.75 (1H, q,  $J$  = 1.3 Hz, CH); <sup>13</sup>C NMR  $\delta$ : 11.8, 12.1, 12.7, 18.9, 25.5, 40.2, 50.8, 60.2, 115.0, 120.1, 124.5, 126.9, 127.7, 147.9, 150.4, 160.4, 167.3; *Anal.* Calcd for C<sub>15</sub>H<sub>22</sub>O<sub>3</sub>: C, 69.84; H, 8.27. Found: C, 70.00; H, 8.51. (ii) *Z*-**14**: pale yellow solids, mp 97 °C; IR  $\nu_{\max}$ : 3462, 1687 cm<sup>-1</sup>; <sup>1</sup>H NMR  $\delta$ : 2.04 (3H, d,  $J$  = 1.3 Hz, Me), 2.21, 2.22, 2.26 (each 3H, s, Me), 2.56-2.60 (2H, m, CH<sub>2</sub>), 2.71-2.76 (2H, m, CH<sub>2</sub>), 3.63, 3.77 (each 3H, s, OMe), 5.78 (1H, q,  $J$  = 1.3

Hz, CH), 7.38 (1H, s, OH, exchangeable);  $^{13}\text{C}$  NMR  $\delta$ : 11.8, 12.3, 12.6, 26.0, 26.5, 34.0, 51.6, 60.3, 115.3, 122.2, 125.4, 128.4, 149.3, 149.8, 161.7, 167.8; *Anal.* Calcd for  $\text{C}_{15}\text{H}_{22}\text{O}_3$ : C, 69.84; H, 8.27. Found: C, 69.77; H, 8.45.

**IOMA of *E*-14 (Table 2, run 1): 6-Methoxy-2-(methoxycarbonylmethyl)-2,5,7,8-tetramethylbenzo[*b*]pyran (15).** To a solution of *E*-14 (50.4 mg, 0.172 mmol) in  $\text{CHCl}_3$  (0.1 mL) was added a solution of TMG in  $\text{CHCl}_3$  (25  $\mu\text{L}/\text{mL}$ , 0.17 mL, 33.9 mmol), and the whole was stirred at rt for 7 h. After evaporation of the solvent, column chromatography of the residue (hexane : AcOEt = 7 : 1) afforded **15** (38.8 mg, 77%) as a colorless oil; IR  $\nu_{\text{max}}$ : 1736  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$ : 1.42 (3H, s, Me), 1.86-1.93, 1.99-2.04 (each 1H, m,  $\text{CH}_2$ ), 2.06, 2.14, 2.18 (each 3H, s, Me), 2.58-2.65 (4H, m,  $\text{CH}_2 \times 2$ ), 3.36, 3.69 (each 3H, s, OMe);  $^{13}\text{C}$  NMR  $\delta$ : 11.6, 12.5, 20.4, 24.7, 31.1, 43.7, 51.5, 60.3, 73.4, 117.1, 123.1, 125.8, 128.0, 146.9, 149.7, 171.1; HREIMS  $m/z$ : 292.1673 (calcd for  $\text{C}_{17}\text{H}_{24}\text{O}_4$ : 292.1674).

**IOMA of *Z*-14 (Table 2, run 4).** A solution of *Z*-14 (50.8 mg, 0.174 mmol) and (-)-Chiba-G (14.6 mg, 0.037 mmol) in  $\text{CHCl}_3$  (0.2 mL) was stirred at rt for 4 days. After evaporation of the solvent, column chromatography of the residue (hexane : AcOEt = 7 : 1) afforded an inseparable 1 : 0.8 mixture of *Z*-14 and **15** (49.8 mg, 98%) as a yellow oil; HPLC (CHIRALCEL OD-H,  $\lambda = 285$  nm, eluent: hexane : *i*-PrOH = 100 : 1, flow rate: 1.1 mL/min):  $t_{\text{R}}$  for a major **15**: 6.79 min (91%),  $t_{\text{R}}$  for a minor **15**: 8.48 min (9%);  $[\alpha]_{\text{D}}^{21} +2.8$  (c 1.0,  $\text{CHCl}_3$ ).

**2-(Formylmethyl)-6-methoxy-2,5,7,8-tetramethylbenzo[*b*]pyran (16).** A 1.04 M solution of DIBAH in hexane (1.0 mL, 1.04 mmol) was added to a solution of the inseparable mixture of *Z*-14 and **15** [0.282 g; calculated to be 0.108 g as **15** (77% ee)] in hexane (16 mL) at  $-7$  °C over 5 min under Ar, and the whole was stirred at the same temperature for 1 h. After successive addition of MeOH (1 mL) and  $\text{H}_2\text{O}$  (20 mL) the mixture was extracted with  $\text{Et}_2\text{O}$  (20 mL  $\times$  3). The combined organic solutions were washed with  $\text{H}_2\text{O}$  (10 mL) and brine (10 mL), dried ( $\text{Na}_2\text{SO}_4$ ), and evaporated. Flash chromatography of the residue (hexane : AcOEt = 20 : 1) afforded **16** (0.044 g, 44% from a calculated **15**) as a colorless oil together with *Z*-14 (0.049 g, 28%); IR  $\nu_{\text{max}}$ : 1721  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$ : 1.40 (3H, s, Me), 1.84, 1.94 (each 1H, ddd,  $J = 13.6, 6.8, 6.8$  Hz,  $\text{CH}_2$ ), 2.08, 2.15, 2.19 (each 3H, s, Me), 2.55 (1H, dd,  $J = 15.0, 3.5$  Hz,  $\text{CH}_2 \times 1/2$ ), 2.61-2.66 (2H, m,  $\text{CH}_2$ ), 2.72 (1H, dd,  $J = 15.0, 2.4$  Hz,  $\text{CH}_2 \times 1/2$ ), 3.64 (3H, s, OMe), 9.93 (1H, dd,  $J = 3.5, 2.4$  Hz, CHO);  $^{13}\text{C}$  NMR  $\delta$ : 11.7, 11.8, 12.5, 20.3, 24.8, 31.9, 52.3, 60.4, 73.5, 117.0, 123.1, 126.0, 128.3, 146.7, 150.0, 202.0; HREIMS  $m/z$ : 262.1568 (calcd for  $\text{C}_{16}\text{H}_{22}\text{O}_3$ : 262.1569).

**(*E*)-Methyl (6-Methoxy-2,5,7,8-tetramethylchroman-2-yl)but-2-enoate (17).** A mixture of **16** (0.041 g, 0.158 mmol) and methyl triphenylphosphonoacetate (0.071 g, 0.214 mmol) in THF (1.75 mL) was

stirred at 45 °C for 2 h. After evaporation of the solvent flash chromatography of the residue (hexane : AcOEt = 20 : 1) afforded **17** (0.032 g, 63%) as a colorless oil; IR  $\nu_{\max}$ : 1723  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$ : 1.27 (3H, s, Me), 1.78, 1.85 (each 1H, ddd,  $J = 19.1, 6.8, 6.8$  Hz,  $\text{CH}_2$ ), 2.09, 2.14, 2.19 (each 3H, s, Me), 2.46 (1H, ddd,  $J = 14.1, 8.1, 1.2$  Hz,  $\text{CH}_2 \times 1/2$ ), 2.53 (1H, ddd,  $J = 14.1, 7.4, 1.2$  Hz,  $\text{CH}_2 \times 1/2$ ), 3.63, 3.74 (each 3H, s, OMe), 5.88 (1H, dt,  $J = 15.7, 1.2$  Hz, CH), 7.05 (1H, dif. t,  $J = 15.7, 7.7$  Hz, CH); HREIMS  $m/z$ : 318.1817 (calcd for  $\text{C}_{19}\text{H}_{26}\text{O}_4$ : 318.1831); HPLC (CHIRALCEL OD-H,  $\lambda = 254$  nm, eluent: hexane :  $i$ PrOH = 150 : 1, flow rate: 1.1 mL/min):  $t_R$  for a major **17**: 12.5 min (84%),  $t_R$  for a minor **17**: 13.6 min (16%);  $[\alpha]_D^{18} -10.8$  ( $c$  1.0,  $\text{CHCl}_3$ ).

## REFERENCES

1. L. J. Machlin, *Vitamin E*, Marcel Dekker, New York, 1980.
2. W. N. Setzer, T. J. Green, R. O. Lawton, D. M. Moriarity, R. B. Bates, S. Caldera, and W. A. Haber, *Planta Med.*, 1995, **61**, 275; J. Z. Deng, D. A. Sun, S. R. Starck, S. M. Hecht, R. L. Cerny, and J. R. Engen, *J. Chem. Soc., Perkin Trans. 1*, 1999, 1147; D. J. Maloney and S. M. Hecht, *Org. Lett.*, 2005, **7**, 4297.
3. Y. Uozumi, K. Kato, and T. Hayashi, *J. Am. Chem. Soc.*, 1997, **119**, 5063.
4. E. Mizuguchi and K. Achiwa, *Chem. Pharm. Bull.*, 1997, **45**, 1209.
5. B. M. Trost and N. Asakawa, *Synthesis*, 1999, 1491; B. M. Trost, H. C. Shen, L. Dong, J.-P. Surivet, and C. Sylvain, *J. Am. Chem. Soc.*, 2004, **126**, 11966.
6. L. F. Tietze, K. M. Sommer, J. Zinngrebe, and F. Stecker, *Angew. Chem. Int. Ed.*, 2005, **44**, 257.
7. M. S. Taylor and E. N. Jacobsen, *Angew. Chem. Int. Ed.*, 2006, **45**, 1520; T. Ishikawa and T. Kumamoto, *Synthesis*, 2006, 737; D. Enders, C. Grondal, and M. R. M. Hüttl, *Angew. Chem. Int. Ed.*, 2007, **46**, 1570, and references cited therein.
8. T. Isobe, K. Fukuda, and T. Ishikawa, *Tetrahedron: Asymmetry*, 1998, **9**, 1729; T. Isobe, K. Fukuda, Y. Araki, and T. Ishikawa, *Chem. Commun.*, 2001, 243; T. Ishikawa, Y. Araki, T. Kumamoto, H. Seki, K. Fukuda, and T. Isobe, *Chem. Commun.*, 2001, 245; K. Hada, T. Watanabe, T. Isobe, and T. Ishikawa, *J. Am. Chem. Soc.*, 2001, **123**, 7705; T. Ishikawa and T. Isobe, *Chem. Eur. J.*, 2002, **8**, 552; Y. Kitani, T. Kumamoto, T. Isobe, K. Fukuda, and T. Ishikawa, *Adv. Synth. Cat.*, 2005, **347**, 1653; D. Wannaporn and T. Ishikawa, *Molecular Diversity*, 2005, **9**, 321; T. Kumamoto, K. Ebine, M. Endo, Y. Araki, Y. Fushimi, I. Miyamoto, T. Ishikawa, T. Isobe, and K. Fukuda, *Heterocycles*, 2005, **66**, 347; T. Haga and T. Ishikawa, *Tetrahedron*, 2005, **61**, 2857; D. Wannaporn and T. Ishikawa, *J. Org. Chem.*, 2005, **70**, 9399; D. Wannaporn, T. Ishikawa, M. Kawahata, and K. Yamaguchi, *J. Org. Chem.*, 2006, **71**, 6600; A. Ryoda, N. Yajima, T. Haga, T. Kumamoto, W. Nakanishi, M. Kawahata, K. Yamaguchi, and T. Ishikawa, *J. Org. Chem.*, 2008, **73**, 133; T. Ishikawa, *Chem. Pharm. Bull.*, 2010, **58**, 1555.

9. N. Saito, A. Ryoda, W. Nakanishi, T. Kumamoto, and T. Ishikawa, *Eur. J. Org. Chem.*, 2008, 2759.
10. E. Sekino, T. Kumamoto, T. Tanaka, T. Ikeda, and T. Ishikawa, *J. Org. Chem.*, 2004, **69**, 2760.
11. C. Ramesh, N. Ravindranath, and B. Das, *J. Org. Chem.*, 2003, **68**, 7101.