

HETEROCYCLES, Vol. 89, No. 1, 2014, pp. 59 - 68. © 2014 The Japan Institute of Heterocyclic Chemistry  
Received, 9th October, 2013, Accepted, 31st October, 2013, Published online, 22nd November, 2013  
DOI: 10.3987/COM-13-12859

## REGIOSELECTIVITY OF THE PALLADIUM-MEDIATED INTRAMOLECULAR COUPLING REACTION OF HIGHLY OXYGENATED PHENYL BENZOATE DERIVATIVES AND APPLICATION TO THE SYNTHESIS OF ALTERTENUOL

Takuya Matsukihira, Shumpei Saga, Yoshikazu Horino, and Hitoshi Abe\*

Graduate School of Science and Engineering, University of Toyama, Gofuku,  
Toyama 930-8555, Japan  
abeh@eng.u-toyama.ac.jp

**Abstract** – The regioselectivity of the intramolecular biaryl coupling reaction of 3',4'-dialkoxyphenyl 2,4-dimethoxybenzoates was investigated using a palladium reagent, and transition state models of the reaction are proposed. As an application of this study, a short-step synthesis of alvertenuol is also performed.


### INTRODUCTION

A palladium-mediated intramolecular biaryl coupling reaction of phenyl benzoate derivatives is a useful transformation for the preparation of the 6*H*-dibenzo[*b,d*]pyran-6-one skeleton.<sup>1</sup> There have been many studies of the total syntheses of natural products possessing this type of heterocyclic system, using the intramolecular biaryl coupling reaction.<sup>2</sup> In some cases, regioselectivity is a crucial concern in the intramolecular biaryl coupling, thus we recently reported several types of reactions involving the issue of regioselectivity.<sup>3</sup> In this report, as a series of our study, regioselectivity is examined in the Pd-mediated intramolecular biaryl coupling reaction of 3',4'-dialkoxyphenyl 2,4-dimethoxybenzoates, and the synthesis of a natural polyphenolic compound, alvertenuol,<sup>4</sup> is also described.

### RESULTS AND DISCUSSION


As the substrates for the investigation, we prepared three kinds of phenyl benzoate derivatives **1a-1c**, which were easily obtained by a simple esterification between the corresponding phenols **2a-2c** and benzoic acid **3**. Initially, the regioselectivity was investigated using Ag<sub>2</sub>CO<sub>3</sub> as the base (Table 1). For the ester **1a**, when Ph<sub>3</sub>P was employed as the ligand, the *ortho*-product **5a** mainly generated. However, a poor regioselectivity for *para*- and *ortho*-products, **4a** and **5a**, was observed under no ligand condition (entries

1 and 2). On the contrary, for the substrates **1b** and **1c**, a dramatic change was observed in the product ratio. Namely, when using  $\text{Ph}_3\text{P}$ , the regioselectivity exhibited 1.4-0.7:1, although when using no ligand, the *para*-isomers **4b** and **4c** were predominantly generated (7.9-8.8:1, entries 3-6).



In contrast to the results above, employing  $\text{K}_2\text{CO}_3$  as the base led to a different outcome for the product ratio. When several phosphine ligands<sup>5</sup> were attempted with the ester **1a**, the regioselectivity was almost identical to the result in entry 1 (entries 7-9). On the other hand, a slightly increased ratio was observed when using no phosphine ligand (entry 10). In the case of both **1b** and **1c**, a moderate selectivity was observed in the range of 1.2-3.6:1 under the condition that the phosphine ligand was employed (entries 11, 13, 14, and 15). When no ligand was used for **1b** and **1c**, the reaction proceeded with the product ratio of 4.4-4.5:1 (entries 12 and 16).

**Table 1.** Reaction of Phenyl Benzoate Derivatives **1a-1c**



entry	substrate	ligand	yield (%)		ratio <sup>b)</sup>	entry	substrate	ligand	yield (%)		ratio <sup>b)</sup>
			4 + 5	4 : 5					4 + 5	4 : 5	
1	<b>1a</b>	$\text{Ph}_3\text{P}$	77	0.2 : 1	7	<b>1a</b>	XPhos <sup>c)</sup>	89	0.3 : 1		
2	<b>1a</b>	none	79	1.8 : 1	8	<b>1a</b>	$\text{Ph}_3\text{P}$	81	0.2 : 1		
3	<b>1b</b>	$\text{Ph}_3\text{P}$	92	1.4 : 1	9	<b>1a</b>	<sup>n</sup> $\text{Bu}_3\text{P}$	77	0.2 : 1		
4	<b>1b</b>	none	89	8.8 : 1	10	<b>1a</b>	none	96	0.5 : 1		
5	<b>1c</b>	$\text{Ph}_3\text{P}$	99	0.7 : 1	11	<b>1b</b>	$\text{Ph}_3\text{P}$	81	2.5 : 1		
6	<b>1c</b>	none	80	7.9 : 1	12	<b>1b</b>	none	84	4.4 : 1		
					13	<b>1c</b>	XPhos <sup>c)</sup>	79	3.6 : 1		
					14	<b>1c</b>	$\text{Ph}_3\text{P}$	89	1.7 : 1		
					15	<b>1c</b>	<sup>n</sup> $\text{Bu}_3\text{P}$	81	1.2 : 1		
					16	<b>1c</b>	none	77	4.5 : 1		


a)  $\text{Ag}_2\text{CO}_3$  was used as the base in entries 1-6.

$\text{K}_2\text{CO}_3$  was used as the base in entries 7-16.

b) Determined by  $^1\text{H-NMR}$  analysis.


c) Ref. 5

For the mechanistic aspect, we would be able to explain the regioselectivity by considering the transition state models.<sup>3i</sup> Based on Echavarren's intermolecular-assisted mechanism,<sup>6</sup> the transition state of substrate **1a** can be illustrated as Figure 1. *A1* and *A2* describe the models in the case of using  $\text{Ag}_2\text{CO}_3$  as the base. Strong affinity between the silver and the iodo atom causes a partial cationic character on the palladium. Therefore, the alkoxy oxygen can coordinate to the palladium atom (*A2*) leading to providing the *ortho*-products **5a-5c**, whereas the steric hindrance, which was generated between the alkoxy group and the ligand on the palladium atom, is also significant (*A2*; L = phosphine). Consequently, the product ratios are controlled by the balance of both the coordinating power and the steric factor. However, when using no phosphine ligand, the tight solvation of the DMA takes place to make the surrounding of the palladium to be bulkier (*A1* and *A2*; L = DMA). Thus, no phosphine ligand condition leads to the severe steric repulsion between the alkoxy group and DMA, as shown in *A2*, so that the ratio of the *para*-products **4a-4c** is increased.




**Figure 1.** Transition State Models under  $\text{Ag}_2\text{CO}_3$  Condition

Contrary to the above, *B1-B3* illustrate the transition state models under the condition using  $\text{K}_2\text{CO}_3$  as the base (Figure 2). In these cases, the cationic property of the palladium atom may not be considered because of the hard character of the potassium ion. The coordinating effect of the alkoxy group to the palladium is not an important factor for the substrates **1b** and **1c**, thus the *para*-isomers **4b** and **4c** are slightly predominant (*B1* and *B2*). Since the coordinating ability of the methylenedioxy group of the substrate **1a** is stronger than **1b** and **1c**, the attractive interaction between the oxygen and the palladium can control and mainly produce the *ortho*-isomer **5a** (*B3*).



**Figure 2.** Transition State Models under  $K_2CO_3$  Condition

Since the coupling product **4c** was regioselectively obtained with the conditions of entry 6 in Table 1, we attempted a further transformation toward the natural product, alternenuol **6**, which was found from *Alternaria tenuis* in 1957.<sup>7</sup> The selective demethylation of **4c** by the Node's method<sup>8</sup> was successful to afford **7**, which was debenzylated by the conventional hydrogenolysis reaction leading to **6**. As an alternative way, partial dealkylation with  $BBr_3$  proceeded effectively to produce **6** in a single step. The spectral data of synthetic **6** were in accordance with the reported data.<sup>4</sup>



**Scheme 1.** Transformation of **4c** into Alternenuol (**6**)

## CONCLUSION

The regioselectivity of the Pd-mediated biaryl coupling reaction of 3',4'-dialkoxyphenyl 2,4-dimethoxybenzoates was intensively investigated. It was found that the combination of the ligand and the base influences the regioselectivity of the reaction. We also demonstrated the synthesis of alvertenuol using the regioselective condition. This method would be utilized for further application to other syntheses of biaryl-type natural products.

## EXPERIMENTAL

### General Information

Melting points were measured using a Yanagimoto micro melting point hot-plate and are uncorrected. The IR spectra were recorded using a SHIMADZU FTIR-8400 spectrophotometer. The NMR spectra were obtained using a JEOL  $\alpha$ -400 (400 MHz) instrument. The chemical shifts are given in  $\delta$  ppm with TMS as an internal standard. The elemental analyses were performed using a Thermo Scientific FlashEA1112 analyzer. Silica gel column chromatography was carried out using a wakogel<sup>®</sup> C-200 or C-400HG (Wako).

### 3,4-Dimethoxyphenol (**2b**)<sup>9</sup>

A mixture of THF (2 mL), 31% H<sub>2</sub>O<sub>2</sub> (0.13 mL), H<sub>3</sub>BO<sub>3</sub> (189 mg, 3.05 mmol), and conc. H<sub>2</sub>SO<sub>4</sub> (0.06 mL) was stirred for 30 min at rt. To the mixture was added a solution of 3,4-dimethoxybenzaldehyde (101 mg, 0.607 mmol) in THF (1 mL). The mixture was stirred for 19 h at rt, then a saturated aqueous NaHCO<sub>3</sub> solution was added. The mixture was extracted with CHCl<sub>3</sub> and the organic layer was washed with brine. After drying with MgSO<sub>4</sub> and evaporation, a residue was subjected to silica gel column chromatography with hexane-AcOEt (4:1) to give **2b** (73.1 mg, 78%) as colorless prisms, mp 83.9-85.2 °C (hexane-CHCl<sub>3</sub>) [lit.,<sup>9</sup> mp 79 °C]. <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ : 6.72 (1H, d,  $J$  = 8.8 Hz, ArH), 6.47 (1H, d,  $J$  = 2.7 Hz, ArH), 6.35 (1H, dd,  $J$  = 8.6, 2.7 Hz, ArH), 5.39 (1H, s, ArOH), 3.81 (3H, s, ArOMe), 3.79 (3H, s, ArOMe).

### 3,4-Dibenzyloxyphenol (**2c**)

A mixture of THF (2 mL), 31% H<sub>2</sub>O<sub>2</sub> (0.07 mL), H<sub>3</sub>BO<sub>3</sub> (100 mg, 1.62 mmol), and conc. H<sub>2</sub>SO<sub>4</sub> (0.032 mL) was stirred for 30 min at rt. To the mixture was added a solution of 3,4-dibenzyloxybenzaldehyde (102 mg, 0.320 mmol) in THF (1.5 mL). The mixture was stirred for 19 h at rt, then a saturated aqueous NaHCO<sub>3</sub> solution was added. The mixture was extracted with CHCl<sub>3</sub> and the organic layer was washed with brine. After drying with MgSO<sub>4</sub> and evaporation, a residue was subjected to silica gel column chromatography with hexane-AcOEt (7:3) to give **2c** (86.8 mg, 88%) as a colorless solid, mp 117.4-118.8

°C (hexane-CHCl<sub>3</sub>). <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 7.44-7.28 (10H, m, ArH), 6.79 (1H, d, *J* = 8.6 Hz, ArH), 6.49 (1H, d, *J* = 3.0 Hz, ArH), 6.29 (1H, dd, *J* = 8.7, 2.8 Hz, ArH), 5.09 (2H, s, ArOCH<sub>2</sub>), 5.07 (2H, s, ArOCH<sub>2</sub>), 4.83 (1H, s, ArOH).

### 3,4-Methylenedioxyphenyl 2-iodo-4,6-dimethoxybenzoate (1a)

Under an N<sub>2</sub> atmosphere, a solution of 2-iodo-4,6-dimethoxybenzoic acid<sup>3d</sup> (104 mg, 0.337 mmol), sesamol (50.7 mg, 0.367 mmol), EDC (90.9 mg, 0.474 mmol), and DMAP (40.7 mg, 0.333 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was stirred for 1 h at rt. The mixture was washed with water and brine, then the organic layer was dried over MgSO<sub>4</sub>. After evaporation, the resulting residue was subjected to silica gel column chromatography with chloroform to give **1** (90.2 mg, 63%). Colorless prisms, mp 110.2-110.9 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). IR (KBr) cm<sup>-1</sup>: 1742, 1593, 1562, 1503, 1478, 1459, 1438, 1408, 1309, 1287, 1243, 1220, 1188, 1169, 1144, 1114, 1097, 1068, 1023, 938, 926, 881, 841, 828, 807, 791, 597. <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 6.95 (1H, d, *J* = 2.2 Hz, ArH), 6.84-6.81 (2H, t, ArH), 6.75 (1H, dd, *J* = 8.4, 2.3 Hz, ArH), 6.48 (1H, d, *J* = 1.9 Hz, ArH) 6.00 (2H, s, ArOCH<sub>2</sub>OAr), 3.85 (3H, s, ArOMe), 3.82 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 166.3, 162.0, 158.2, 148.1, 145.7, 145.2, 122.4, 115.4, 114.2, 108.1, 103.9, 101.8, 99.1, 92.9, 56.2, 55.9. *Anal.* Calcd for C<sub>16</sub>H<sub>13</sub>IO<sub>6</sub>: C, 44.88; H, 3.06. Found: C, 44.74; H, 2.94.

### 3,4-Dimethoxyphenyl 2-iodo-4,6-dimethoxybenzoate (1b)

In the same procedure mentioned above using **2b** (52.8 mg, 0.343 mmol), **1b** (63.4 mg, 44%) was obtained. Colorless prisms, mp 80.2-80.5 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). IR (KBr) cm<sup>-1</sup>: 1749, 1598, 1562, 1510, 1455, 1255, 1228, 1182, 1151, 1124, 1076, 1030. <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 6.95 (1H, d, *J* = 2.2 Hz, ArH), 6.87 (2H, d, *J* = 1.5 Hz, ArH), 6.84 (1H, t, *J* = 1.5 Hz), 6.47 (1H, d, *J* = 2.2 Hz, ArH) 3.88 (3H, s, ArOMe), 3.87 (3H, s, ArOMe), 3.83 (3H, s, ArOMe), 3.80 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 166.3, 161.9, 158.1, 149.5, 147.2, 144.6, 122.5, 115.4, 113.0, 111.3, 105.9, 99.0, 92.9, 56.3, 56.2, 56.1, 55.8. *Anal.* Calcd for C<sub>17</sub>H<sub>17</sub>IO<sub>6</sub>: C, 45.96; H, 3.86. Found: C, 45.84; H, 3.72.

### 3,4-Dibenzyloxyphenyl 2-iodo-4,6-dimethoxybenzoate (1c)

In the same procedure mentioned above using **2c** (4.04 g, 13.2 mmol), **1c** (4.87 g, 63%) was obtained. Colorless solid, mp 101.4-102.0 °C (Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub>). IR (KBr) cm<sup>-1</sup>: 1751, 1603, 1560, 1510, 1454, 1435, 1421, 1400, 1387, 1331, 1315, 1277, 1250, 1236, 1215, 1184, 1153, 1119, 1074, 1030, 1001, 989, 922, 883, 818, 804, 787, 770, 750, 731, 702, 594. <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 7.47-7.29 (10H, m, ArH), 6.97-6.93 (3H, m, ArH), 6.84 (1H, dd, *J* = 8.7, 2.6 Hz, ArH), 6.47 (1H, d, *J* = 2.2 Hz, ArH), 5.18 (2H, s,

ArOCH<sub>2</sub>), 5.16 (2H, s, ArOCH<sub>2</sub>), 3.82 (3H, s, ArOMe), 3.81 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 166.1, 161.9, 158.2, 149.6, 147.0, 145.1, 137.3, 137.0, 128.6, 128.6, 128.0, 127.9, 127.5, 127.5, 122.5, 115.6, 115.4, 114.0, 109.1, 99.1, 93.0, 72.0, 71.5, 56.2, 55.8. *Anal.* Calcd for C<sub>29</sub>H<sub>25</sub>IO<sub>6</sub>: C, 58.40; H, 4.23. Found: C, 58.62; H, 4.15.

### Typical Procedure for the Coupling Reaction (Table 1)

Under an N<sub>2</sub> atmosphere, a mixture of **1a** (100 mg, 0.234 mmol), Pd(OAc)<sub>2</sub> (0.023 mmol), ligand (0.047 mmol), base (0.023 mmol), and DMA (7 mL) was heated under reflux for 1 h with stirring. Unsolved materials were filtered off, then water was added to the filtrate. After extraction with CH<sub>2</sub>Cl<sub>2</sub>, the organic layer was washed with brine and dried over MgSO<sub>4</sub>. Evaporation of the solvent gave a residue which was subjected to silica gel column chromatography with hexane-AcOEt-CH<sub>2</sub>Cl<sub>2</sub> (1:1:3).

### 2,3-Methylenedioxy-7,9-dimethoxy-6H-benzo[*c*]chromen-6-one (**4a**)

Colorless needles, mp 240.5-241.0 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 7.25 (1H, s, ArH), 6.83 (1H, d, *J* = 2.2 Hz, ArH), 6.75 (1H, s, ArH), 6.48 (1H, d, *J* = 2.2 Hz, ArH), 6.04 (2H, s, ArOCH<sub>2</sub>OAr), 3.98 (3H, s, ArOMe), 3.96 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 165.5, 164.2, 158.1, 150.0, 148.1, 145.0, 139.7, 111.1, 102.9, 102.2, 101.2, 98.6, 98.3, 96.8, 56.5, 55.8. IR (KBr) cm<sup>-1</sup>: 1719, 1603, 1574, 1504, 1472, 1454, 1433, 1381, 1344, 1254, 1211, 1180, 1146, 1092, 1063, 1030, 1005, 827. *Anal.* Calcd for C<sub>16</sub>H<sub>12</sub>O<sub>6</sub>: C, 64.00; H, 4.03. Found: C, 63.85; H, 3.84.

### 1,2-Methylenedioxy-7,9-dimethoxy-6H-benzo[*c*]chromen-6-one (**5a**)

Colorless needles, mp 254.8-256.5 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 7.58 (1H, d, *J* = 2.2 Hz, ArH), 6.91 (1H, d, *J* = 8.6 Hz, ArH), 6.80 (1H, d, *J* = 8.8 Hz, ArH), 6.56 (1H, d, *J* = 2.4 Hz, ArH), 6.16 (2H, s, ArOCH<sub>2</sub>OAr), 4.00 (3H, s, ArOMe), 3.95 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 165.5, 163.8, 157.9, 146.8, 144.0, 143.7, 137.2, 109.8, 109.0, 105.3, 103.4, 102.3, 101.8, 99.4, 56.5, 55.8. IR (KBr) cm<sup>-1</sup>: 1722, 1641, 1612, 1597, 1574, 1504, 1489, 1456, 1445, 1418, 1346, 1265, 1246, 1205, 1165, 1080, 1057, 1043, 1007, 986, 920, 839, 916. *Anal.* Calcd for C<sub>16</sub>H<sub>12</sub>O<sub>6</sub>: C, 64.00; H, 4.03. Found: C, 63.92; H, 3.96.

### 2,3,7,9-Tetramethoxy-6H-benzo[*c*]chromen-6-one (**4b**)

Colorless solid, mp 180.8-181.7 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 7.21 (1H, s, ArH), 6.87 (1H, d, *J* = 2.2 Hz, ArH), 6.77 (1H, s, ArH), 6.46 (1H, d, *J* = 2.2 Hz, ArH), 3.98 (3H, s, ArOMe), 3.98 (3H, s, ArOMe), 3.97 (3H, s, ArOMe), 3.91 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 165.1, 163.8, 157.9, 151.4, 146.5, 145.7, 139.0, 109.3, 104.0, 102.5, 99.8, 97.5, 96.2, 56.3, 56.0, 56.0, 55.5. IR (KBr) cm<sup>-1</sup>: 3576, 1686, 1618, 1597, 1570, 1520, 1458, 1439, 1425, 1408, 1273, 1238, 1211, 1198, 1173,

1159, 1123, 1070, 995, 824. *Anal.* Calcd for C<sub>17</sub>H<sub>16</sub>O<sub>6</sub>: C, 64.55; H, 5.10. Found: C, 64.33; H, 4.98.

### 1,2,7,9-Tetramethoxy-6*H*-benzo[*c*]chromen-6-one (5b)

Colorless solid, mp 202.2-203.5 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 8.33 (1H, d, *J* = 2.2 Hz, ArH), 7.07 (2H, s, ArH), 6.61 (1H, d, *J* = 2.4 Hz, ArH), 3.99 (3H, s, ArOMe), 3.96 (3H, s, ArOMe), 3.92 (3H, s, ArOMe), 3.91 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 165.4, 163.7, 157.9, 149.5, 147.3, 146.4, 138.5, 114.7, 112.7, 112.4, 104.1, 102.3, 99.6, 60.4, 56.7, 56.5, 55.6. IR (KBr) cm<sup>-1</sup>: 1724, 1601, 1587, 1570, 1497, 1474, 1456, 1443, 1414, 1337, 1298, 1267, 1246, 1229, 1213, 1204, 1163, 1090, 1072, 1059, 1024, 962, 837, 804. *Anal.* Calcd for C<sub>17</sub>H<sub>16</sub>O<sub>6</sub>: C, 64.55; H, 5.10. Found: C, 64.45; H, 5.05.

### 2,3-Dibenzyloxy-7,9-dimethoxy-6*H*-benzo[*c*]chromen-6-one (4c)

Colorless solid, mp 166.5-167.3 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 7.49-7.30 (11H, m, ArH), 6.81 (1H, s, ArH), 6.72 (1H, d, *J* = 2.2 Hz, ArH), 6.42 (1H, d, *J* = 2.2 Hz, ArH), 5.20 (2H, s, ArOCH<sub>2</sub>), 5.18 (2H, s, ArOCH<sub>2</sub>), 3.95 (3H, s, ArOMe), 3.92 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 165.4, 164.2, 158.2, 152.0, 147.4, 145.4, 139.4, 137.2, 136.2, 128.8, 128.7, 128.3, 128.1, 127.6, 127.3, 110.2, 109.9, 103.0, 102.5, 98.1, 96.6, 72.8, 71.0, 56.4, 55.7. IR (KBr) cm<sup>-1</sup>: 1738, 1614, 1593, 1568, 1518, 1499, 1448, 1431, 1391, 1333, 1244, 1211, 1196, 1159, 1126, 1061, 1020, 1003, 924, 839, 824, 733, 694. *Anal.* Calcd for C<sub>29</sub>H<sub>24</sub>O<sub>6</sub>: C, 74.35; H, 5.16. Found: C, 74.47; H, 5.22.

### 1,2-Dibenzyloxy-7,9-dimethoxy-6*H*-benzo[*c*]chromen-6-one (5c)

Colorless solid, mp 184.4-186.1 °C (hexane-CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ: 8.29 (1H, d, *J* = 2.4 Hz, ArH), 7.47-7.31 (10H, m, ArH), 7.14 (1H, d, *J* = 9.0 Hz, ArH), 7.06 (1H, d, *J* = 9.0 Hz, ArH), 6.55 (1H, d, *J* = 2.2 Hz, ArH), 5.16 (2H, s, ArOCH<sub>2</sub>), 5.08 (2H, s, ArOCH<sub>2</sub>), 3.96 (3H, s, ArOMe), 3.43 (3H, s, ArOMe). <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ: 165.4, 163.5, 157.9, 148.7, 146.8, 138.7, 137.0, 136.7, 128.7, 128.6, 128.5, 128.4, 128.3, 127.8, 117.0, 113.2, 112.6, 104.1, 102.2, 100.2, 75.2, 72.1, 56.4, 55.3. IR (KBr) cm<sup>-1</sup>: 1742, 1601, 1572, 1501, 1474, 1456, 1443, 1427, 1371, 1337, 1290, 1261, 1252, 1215, 1196, 1167, 1049, 1022, 959, 800, 752, 731, 696. *Anal.* Calcd for C<sub>29</sub>H<sub>24</sub>O<sub>6</sub>: C, 74.35; H, 5.16. Found: C, 74.26; H, 5.13.

### Altertenuol (2,3,7-Trihydroxy-9-methoxy-6*H*-benzo[*c*]chromen-6-one) (6)

Under N<sub>2</sub> atmosphere, to a solution of **4c** (50.1 mg, 0.107 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3.5 mL) was added BBr<sub>3</sub> (1M CH<sub>2</sub>Cl<sub>2</sub> solution, 0.320 mL, 0.320 mmol) at 0 °C. The mixture was stirred for 1 h at rt, and quenched with water. After extraction with AcOEt, the organic layer was washed with brine and dried over MgSO<sub>4</sub> to give a residue which was subjected to silica gel column chromatography with AcOEt. A colorless solid of **1** (27.9 mg, 95%) was obtained, mp 283.9-285.8 °C (acetone) [lit.,<sup>7</sup> mp 284-285 °C]. <sup>1</sup>H-NMR (400 MHz, CD<sub>3</sub>OD) δ: 7.38 (1H, s, ArH), 6.94 (1H, d, *J* = 2.4 Hz, ArH), 6.74 (1H, s, ArH), 6.50 (1H, d, *J* =



2.2 Hz, ArH), 3.93 (3H, s, ArOMe).  $^{13}\text{C}$ -NMR (100 MHz, DMSO- $d_6$ )  $\delta$ : 166.8, 165.1, 163.8, 149.3, 144.4, 143.8, 137.6, 109.1, 109.0, 103.5, 100.0, 98.6, 98.0, 56.1. IR (KBr)  $\text{cm}^{-1}$ : 3535, 3285, 3105, 1647, 1607, 1568, 1528, 1510, 1456, 1431, 1279, 1227, 1202, 1165. *Anal.* Calcd for  $\text{C}_{14}\text{H}_{10}\text{O}_6$ : C, 61.32; H, 3.68. Found: C, 61.26; H, 3.69.

### 2,3-Dibenzyloxy-7-hydroxy-9-methoxy-6H-benzo[c]chromen-6-one (7)<sup>4a</sup>

Under  $\text{N}_2$  atmosphere, to a solution of **4c** (102 mg, 0.218 mmol) in a mixed solvent of  $\text{CH}_2\text{Cl}_2$  (4.7 mL) and MeCN (1.4 mL) was added  $\text{AlCl}_3$  (69.6 mg, 0.522 mmol) and NaI (83.3 mg, 0.556 mmol) at 0 °C. The mixture was stirred for 4.5 h at rt, and quenched with water. After extraction with  $\text{CH}_2\text{Cl}_2$ , the organic layer was washed with brine and dried over  $\text{MgSO}_4$  to give a residue which was subjected to silica gel column chromatography with hexane-AcOEt (4:3). Colorless solid of **7** (69.9 mg, 71%) was obtained, mp 211.3-212.5 °C (AcOEt) [lit.,<sup>4a</sup> mp 199-200 °C].  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 11.5 (1H, s, ArOH), 7.49-7.32 (11H, m, ArH), 6.88 (1H, s, ArH), 6.75 (1H, d,  $J = 2.2$  Hz, ArH), 6.50 (1H, d,  $J = 2.2$  Hz, ArH), 5.22 (4H, s,  $\text{ArOCH}_2$ ), 3.91 (3H, s, ArOMe).

### Conversion of 7 into 6

Under  $\text{H}_2$  atmosphere, a mixture of **7** (53.6 mg, 0.118 mmol), 10% Pd/C (12.1 mg), and THF (4 mL) was vigorously stirred for 19 h at rt. After filtration and washing the filter paper with  $\text{CHCl}_3$ -MeOH (10:1), the filtrate was concentrated. An obtained residue was purified by silica gel column chromatography with AcOEt to give **6** (31.6 mg, 97%).

### ACKNOWLEDGMENT

A part of this work was financially supported by JSPS KAKENHI Grant No. 22590003 and the Tamura Science and Technology Foundation.

### REFERENCES AND NOTES

- (a) D. E. Ames and A. Opalko, *Tetrahedron*, 1984, **40**, 1919; For reviews, see: (b) G. Bringmann, M. Breuning, and S. Tasler, *Synthesis*, 1999, 525; (c) J. Hassan, M. Sévignon, C. Gozzi, E. Schulz, and M. Lemaire, *Chem. Rev.*, 2002, **102**, 1359; (d) G. Bringmann, A. J. P. Mortimer, P. A. Keller, M. J. Gresser, J. Garner, and M. Breuning, *Angew. Chem. Int. Ed.*, 2005, **44**, 5384; (e) D. Alberico, M. E. Scott, and M. Lautens, *Chem. Rev.*, 2007, **107**, 174; (f) G. Bringmann, T. Gulder, T. A. M. Gulder, and M. Breuning, *Chem. Rev.*, 2011, **111**, 563; (g) J. Yamaguchi, A. D. Yamaguchi, and K. Itami, *Angew. Chem. Int. Ed.*, 2012, **51**, 8960.
- (a) T. Matsumoto, T. Hosoya, and K. Suzuki, *J. Am. Chem. Soc.*, 1992, **114**, 3568; (b) T. Hosoya, E.

- Takashiro, T. Matsumoto, and K. Suzuki, *J. Am. Chem. Soc.*, 1994, **116**, 1004; (c) T. Harayama, H. Yasuda, T. Akiyama, Y. Takeuchi, and H. Abe, *Chem. Pharm. Bull.*, 2000, **48**, 861; (d) G. Bringmann and D. Menche, *Acc. Chem. Res.*, 2001, **34**, 615; (e) G. Bringmann, M. Breuning, R.-M. Pfeifer, W. A. Schenk, K. Kamikawa, and M. Uemura, *J. Organomet. Chem.*, 2002, **661**, 31; (f) G. Bringmann, S. Tasler, R.-M. Pfeifer, and M. Breuning, *J. Organomet. Chem.*, 2002, **661**, 49; (g) G. A. Molander, K. M. George, and L. G. Monovich, *J. Org. Chem.*, 2003, **68**, 9533; (h) H. Abe, S. Takeda, T. Fujita, K. Nishioka, Y. Takeuchi, and T. Harayama, *Tetrahedron Lett.*, 2004, **45**, 2327; (i) H. Abe, K. Nishioka, S. Takeda, M. Arai, Y. Takeuchi, and T. Harayama, *Tetrahedron Lett.*, 2005, **46**, 3197; (j) S. Takeda, H. Abe, Y. Takeuchi, and T. Harayama, *Tetrahedron*, 2007, **63**, 396; (k) H. Abe, Y. Sahara, Y. Matsuzaki, Y. Takeuchi, and H. Abe, *Tetrahedron Lett.*, 2008, **49**, 605; (l) H. Abe and T. Harayama, *Heterocycles*, 2008, **75**, 1305; (m) K. Shioe, Y. Takeuchi, T. Harayama, and H. Abe, *Chem. Pharm. Bull.*, 2010, **58**, 435; (n) K. Shioe, Y. Sahara, Y. Horino, T. Harayama, Y. Takeuchi, and H. Abe, *Tetrahedron*, 2011, **67**, 1960.
3. (a) T. Harayama, Y. Kawata, C. Nagura, T. Sato, T. Miyagoe, H. Abe, and Y. Takeuchi, *Tetrahedron Lett.*, 2005, **46**, 6091; (b) T. Harayama, A. Hori, H. Abe, and Y. Takeuchi, *Heterocycles*, 2006, **67**, 385; (c) H. Nishioka, C. Nagura, H. Abe, Y. Takeuchi, and T. Harayama, *Heterocycles*, 2006, **70**, 549; (d) H. Abe, T. Fukumoto, Y. Takeuchi, and T. Harayama, *Heterocycles*, 2007, **74**, 265; (e) T. Harayama, M. Asai, T. Miyagoe, H. Abe, Y. Takeuchi, A. Yamaguchi, and S. Fujii, *Heterocycles*, 2010, **81**, 1881; (f) T. Harayama, C. Nagura, T. Miyagoe, Y. Kawata, H. Abe, and Y. Takeuchi, *Heterocycles*, 2010, **81**, 2609; (g) H. Abe, T. Fukumoto, Y. Horino, T. Harayama, and Y. Takeuchi, *Heterocycles*, 2010, **82**, 851; (h) H. Abe, T. Matsukihira, T. Fukumoto, Y. Horino, Y. Takeuchi, and T. Harayama, *Heterocycles*, 2012, **84**, 323; (i) K. Maeda, T. Matsukihira, S. Saga, Y. Takeuchi, T. Harayama, Y. Horino, and H. Abe, *Heterocycles*, 2014, **88**, 621.
  4. (a) K. Kuramochi, K. Fukudome, I. Kuriyama, T. Takeuchi, Y. Sato, S. Kamisuki, K. Tsubaki, S. Sugawara, H. Yoshida, and Y. Mizushina, *Bioorg. Med. Chem.*, 2009, **17**, 7227; (b) G. Nemecek, J. Cudaj, and J. Podkech, *Eur. J. Org. Chem.*, 2012, 3863.
  5. For Xphos, see: K. Billingsley and S. L. Buchwald, *J. Am. Chem. Soc.*, 2007, **129**, 3358.
  6. (a) D. Garcia-Cuadrado, A. A. C. Braga, F. Maseras, and A. M. Echavarren, *J. Am. Chem. Soc.*, 2006, **128**, 1066; (b) D. Garcia-Cuadrado, P. de Mendoza, A. A. C. Braga, F. Maseras, and A. M. Echavarren, *J. Am. Chem. Soc.*, 2007, **129**, 6880.
  7. (a) T. Rosett, R. H. Sankhala, C. E. Stickings, M. E. U. Taylor, and R. Thomas, *Biochem. J.*, 1957, **67**, 390; (b) R. Thomas, *Biochem. J.*, 1961, **80**, 234.
  8. M. Node, T. Kajimoto, K. Nishide, E. Fujita, and K. Fuji, *Bull. Inst. Chem. Res., Kyoto Univ.*, 1992, **70**, 308.
  9. A. Roy, K. R. Reddy, P. K. Mohanta, H. Ila, and H. Junjappa, *Synth. Commun.*, 1999, **29**, 3781.