

Supporting information

A new iridoid glycoside from *Santisukia pagetii*

Poolsak Sahakitpichan,¹ Nitirat Chimnoi,¹ Chutima Srinroch,¹ Chaleaw Petchthong,²
Somsak Ruchirawat,¹ and Tripetch Kanchanapoom^{1,3*}

¹ Chulabhorn Research Institute, Kamphaeng Phet 6, Talat Bang Khen, Lak Si, Bangkok 10210, Thailand.

² Faculty of Science and Technology, Kanchanaburi Rajabhat University, Kanchanaburi 71190, Thailand.

³ Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen 40002, Thailand.

*Corresponding author: E-mail: trikan@kku.ac.th

Contents

The ^1H NMR spectrum of compound 1 in CD_3OD	Figure S1
The ^{13}C NMR spectrum of compound 1 in CD_3OD	Figure S2
The COSY spectrum of compound 1 in CD_3OD	Figure S3
The HSQC spectrum of compound 1 in CD_3OD	Figure S4
The HMBC spectrum of compound 1 in CD_3OD	Figure S5




Figure S1 The ^1H NMR spectrum of compound **1** in CD_3OD




Figure S2 The ^{13}C NMR spectrum of compound **1** in CD_3OD




Figure S3 The COSY spectrum of compound **1** in CD₃OD




Figure S4 The HSQC spectrum of compound **1** in CD_3OD




Figure S5 The HMBC spectrum of compound **1** in CD₃OD