

**MACROCARQUINOID D, NEW MEROTERPENOID FROM BROWN
ALGA, *SARGASSUM MACROCARPUM***

Seiya Shinoda¹, Shin-ichiro Kurimoto², Takaaki Kubota³, and Mitsuhiro Sekiguchi^{1*}

¹ Graduate School of Bioresources and Environmental Sciences, Ishikawa Prefectural University, 1-308 Suematsu Nonoichi-shi, Ishikawa 921-8836, Japan. ² School of Pharmacy, Showa University, 1-5-8 Hatanodai Shinagawa-ku, Tokyo 142-8555, Japan. ³ Faculty of Medicine, Dentistry, and Pharmaceutical Sciences, Okayama University, 1-1-1 Tsushima-naka, Kita-ku, Okayama 700-8530, Japan.

Content

Fig. S1: ^1H NMR spectrum of **1** in CDCl_3

Fig. S2: ^{13}C NMR spectrum of **1** in CDCl_3

Fig. S3: ^1H - ^1H COSY spectrum of **1** in CDCl_3

Fig. S4: HSQC spectrum of **1** in CDCl_3

Fig. S5: HMBC spectrum of **1** in CDCl_3

Fig. S6: NOESY spectrum of **1** in CDCl_3




Fig. S1: ^1H NMR spectrum of **1** in CDCl_3




Fig. S2: ^{13}C NMR spectrum of **1** in CDCl_3




Fig. S3: ^1H - ^1H COSY spectrum of **1** in CDCl_3




Fig. S4: HSQC spectrum of **1** in CDCl_3




Fig. S5: HMBC spectrum of **1** in CDCl_3




Fig. S6: NOESY spectrum of **1** in CDCl_3